arXiv:1207.3562v1 [hep-lat] 16 Jul 2012

ADP-12-30/T797

Baryon resonances and hadronic interactions in a finite volume

J. M. M. Hall,[1] A. C.-P. Hsu,[1] D. B. Leinweber,[1] A. W. Thomas,[1,2] and R. D. Young[1,2]

[1]*Special Research Centre for the Subatomic Structure of Matter (CSSM),*
School of Chemistry and Physics, University of Adelaide 5005, Australia
[2]*ARC Centre of Excellence for Particle Physics at the Terascale,*
School of Chemistry and Physics, University of Adelaide 5005, Australia

The connection between the asymptotic states of the S-matrix and the energy spectrum in a finite volume is investigated in the context of a $\Delta N\pi$ interaction model that can be solved exactly. In particular, Lüscher's method, which proposes a definite relationship between the elastic scattering phase shift and the finite-volume energy levels, is shown to exhibit a deviation from the correct resonance position that behaves approximately like $1/L$ in the range of volumes of interest. A new method is proposed, whereby the energy levels from a lattice calculation are used to determine the parameters of the model. The phase shift may then be evaluated using an infinite-volume t-matrix. The model-dependence of the method is investigated. At box sizes that are currently considered quite large, model-dependent effects are estimated to be an order of magnitude smaller than the deviations associated with Lüscher's method. The extraction of the resonance position is also shown to be relatively independent of L.

PACS numbers: 12.38.Gc 12.38.Aw 12.39.Fe

Recent advances in lattice QCD have demonstrated the ability to extract the low-lying tower of excitation energies reliably in baryon correlators [1–3]. Nevertheless, the identification of resonance parameters (e.g. masses, widths, etc.) from multi-hadron interactions remains an ongoing challenge of current and active research [4–14]. In particular, the calculation of the discrete energy levels of a hadronic system on a finite-volume lattice needs to be related to the continuous spectra measured by scattering experiments. One approach, Lüscher's method [15], allows such a relation to be made by geometrically matching the asymptotic behaviour of the S-matrix onto the energy spectrum of the toroidal topology of the lattice. The standard implementation of Lüscher's method neglects corrections to energy levels that are exponentially suppressed in the lattice volume. In this Letter, an exactly solvable model is used to investigate the role of these terms in the extraction of resonance parameters.

Lüscher's method entails a fixed relationship between the scattering phase shift δ, and the rth energy level in a finite volume (in keeping with Lüscher's notation, r is an integer):

$$r\,\pi - \delta(k_r) = \phi(q_r). \tag{1}$$

$q_r = k_r L/(2\pi)$ are the (dimensionless) lattice momenta, and $\phi(q)$ is a geometric function related to the three-dimensional Zeta function, defined by:

$$\phi(q) = -\arctan\left(\frac{\pi^{3/2} q}{\mathcal{Z}_{00}(1,q^2)}\right), \tag{2}$$

$$\mathcal{Z}_{00}(1,q^2) = \frac{1}{\sqrt{4\pi}} \sum_{\vec{n}\in\mathbb{Z}^3} \frac{1}{\vec{n}^2 - q^2}. \tag{3}$$

A simple minimal subtraction scheme has been chosen to regulate the divergence in the Zeta function.

Lüscher's method will be tested in the context of a well-studied model describing the decay of the Δ-baryon into $N\pi$; an archetypal example of a baryon resonance. It will be shown that Lüscher's method entails a deviation from the infinite-volume resonance position of order $\sim 1/L$, for moderate and even large volumes. A new method is proposed, in which energy levels from the lattice are fit to those of the model, and the phase shift then calculated using the infinite-volume t-matrix.

In the following finite-volume heavy-baryon model, a matrix Hamiltonian, $H = H_0 + H_I$, is defined to include the interaction between the Δ-baryon and the pion-nucleon system. The rows and columns of H represent the momentum values available to the pion ($k_1 = 2\pi/L, \ldots$). In this model, the $N\pi$ states are only coupled to the Δ-baryon. The free Hamiltonian, H_0, takes the form:

$$H_0 = \begin{pmatrix} \Delta_0 & 0 & 0 & \cdots \\ 0 & \omega_\pi(k_1) & 0 & \cdots \\ 0 & 0 & \omega_\pi(k_2) & \\ \vdots & \vdots & & \ddots \end{pmatrix}, \tag{4}$$

where $\omega_\pi(k_n) \equiv \sqrt{k_n^2 + m_\pi^2}$, and the momenta at finite volume satisfies the condition: $k_n^2 = \left(\frac{2\pi}{L}\right)^2 n$, for a squared integer $n \equiv n_x^2 + n_y^2 + n_z^2$. The matrix H_0 has non-zero diagonal entries representing the non-interacting energy of a pion at certain values of momentum, and the nucleon recoil vanishes in the heavy-baryon limit. Note that the cubic symmetry of the system means that the dimension of H is significantly smaller; each non-zero n occurs only once. The first element of the H_0 includes a bare resonance mass relative to the nucleon mass, Δ_0, which is chosen so that the phase shift passes through 90° at 292 MeV in the infinite-volume limit.

The interacting Hamiltonian, H_I, contains couplings, $g_{\Delta N}$, in the top row and left-most column:

$$H_I = \begin{pmatrix} 0 & g^{\rm fin}_{\Delta N}(k_1) & g^{\rm fin}_{\Delta N}(k_2) & \cdots \\ g^{\rm fin}_{\Delta N}(k_1) & 0 & 0 & \cdots \\ g^{\rm fin}_{\Delta N}(k_2) & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}. \tag{5}$$

The $\Delta \leftrightarrow N\pi$ coupling, $g^{\rm fin}_{\Delta N}(k_n)$, is defined in terms of known phenomenological parameters $f_\pi = 92.4$ MeV, and

$\mathcal{C} = -1.52$, as derived from the SU(6) flavour-symmetry relation, and must also contain a normalization factor $\sqrt{C_3(n)}$, representing the number of ways of summing the squares of three integers to equal n:

$$g^{\rm fin}_{\Delta N}(k_n) = \sqrt{\frac{C_3(n)}{4\pi}} \left(\frac{2\pi}{L}\right)^{3/2} g_{\Delta N}(k_n) \tag{6}$$

$$= \sqrt{\chi_\Delta \frac{C_3(n)}{2\pi^2}} \left(\frac{2\pi}{L}\right)^{3/2} \frac{k_n\, u(k_n)}{\sqrt{\omega_\pi(k_n)}}, \tag{7}$$

$$\text{for} \quad \chi_\Delta = \frac{3}{32\pi f_\pi^2} \frac{2}{9} \mathcal{C}^2, \tag{8}$$

$$\text{and} \quad u(k_n) = \left(1 + \frac{k_n^2}{\Lambda^2}\right)^{-2}. \tag{9}$$

Note that a regulator, $u(k)$, has been introduced into the coupling term. This will allow the form of the energy levels to be related to the regularized one-pion loop integral occurring in the t-matrix, as shown below. (For detailed discussions of finite-range regularization, see Refs. [16–20]). For the purposes of this model, a dipole regulator with $\Lambda = 0.8$ GeV is chosen. It will be shown, however, that the form of the regulator has little impact on the consistency of the final result: extracting the correct resonance position.

One extension of this model would be to include a direct $N\pi \leftrightarrow N\pi$ interaction, which could be viewed as an approximation to the famous Chew-Low interaction. For example, one could add additional terms in the interaction matrix H_I:

$$g^{\rm fin}_{CL}(k_i, k_j) = -g_{CL}(k_i) g_{CL}(k_j) \frac{1}{\sqrt{\omega_\pi(k_i)\omega_\pi(k_j)}}. \tag{10}$$

In the Cloudy Bag Model [21, 22], The couplings g_{CL} and $g^{\rm fin}_{CL}$ are related by the equation: $g_{CL}(k_i) = \kappa g^{\rm fin}_{\Delta N}(k_i)$, with a ratio, $\kappa = \sqrt{25/18}$, estimated from SU(6) symmetry. In the case of the t-matrix, the Lippmann-Schwinger equation with a rank-2 separable potential can be solved as per Mongan [23]. Suffice to say, this additional interaction will be considered only in the final analysis, as a test of model-independence.

Firstly, consider only the $\Delta \leftrightarrow N\pi$ interaction. The eigenvalues of the matrix H correspond to the different energy levels in a finite volume, and may be calculated for a range of L values. The ten lowest energy levels as a function of L are shown in Fig. 1, with the non-interacting energies shown as dotted lines. It will be useful to have a reliable method for extracting phase shifts in this region, even if larger box sizes were feasible in lattice QCD.

In the case of this model, the phase shift can, in fact, be calculated exactly, thus providing a suitable test for the ability Lüscher's method to reproduce the correct phase shift. The solution to the eigenvalue problem for H takes the following form (provided $\omega_\pi(k_n) \neq \lambda$):

$$\lambda = \Delta_0 - \sum_{n=1}^{\infty} \frac{\left(g^{\rm fin}_{\Delta N}(k_n)\right)^2}{\omega_\pi(k_n)\, - \lambda}, \tag{11}$$

which mirrors the t-matrix calculation for elastic $N\pi$-scattering with a Δ-baryon intermediate. The relationship between the phase shift and the on-shell quantity, $T = t(k,k;E^+)$, is:



FIG. 1. (color online). The ten lowest energy levels from the $\Delta N\pi$ model described in Eqs. (4)−(9) (solid lines), and the corresponding non-interacting energies (dotted lines).

$$T = \frac{g^2_{\Delta N}(k)}{\omega_\pi(k) - \Delta_0 - \Sigma_{\Delta N}(k)} = -\frac{1}{\pi k\, \omega_\pi(k)}\, e^{i\delta(k)} \sin\delta(k). \tag{12}$$

The bare resonance mass, Δ_0, may be tuned so that the final resonance energy matches the physical ΔN mass-splitting: 292 MeV, and consequently fed into Eq. (4).

The $\Delta \leftrightarrow N\pi$ one-pion loop integral in a finite volume is equal to the sum in Eq. (11), matching the effective field theory calculation in the weak coupling limit, $E_\Delta \to \Delta_0$:

$$E = \Delta_0 - \frac{\chi_\Delta}{2\pi^2} \left(\frac{2\pi}{L}\right)^3 \sum_{n=1}^{\infty} C_3(n) \frac{k_n^2 u^2(k_n)}{\omega_\pi(k_n)[\omega_\pi(k_n)\, - E_\Delta]}. \tag{13}$$

At infinite volume, the real part of the loop integral takes the form:

$$\mathrm{Re}\,\Sigma_{\Delta N}(k) = \mathcal{P}\int_0^\infty \mathrm{d}k' \frac{k'^{\,2} g^2_{\Delta N}(k'\,)}{\omega_\pi(k) - \omega_\pi(k'\,)} \tag{14}$$

$$= \chi_\Delta \frac{2}{\pi} \mathcal{P}\int_0^\infty \mathrm{d}k' \frac{k'^{\,4}\, u^2(k'\,)}{\omega_\pi(k'\,)\,[\omega_\pi(k) - \omega_\pi(k'\,)]}, \tag{15}$$

where $\mathcal{P}$ indicates that a principal value integral must be performed. By solving Eq. (12), the phase shift, δ, may be plotted as a function of energy, $E = \omega_\pi(k)$, to obtain the Breit-Wigner-like curve shown in Fig. 2.

In order to test Lüscher's formula for the $\Delta N\pi$ model, the phase shift is calculated from the eigenvalues obtained from H using Eq. (1). This phase shift can be compared to the infinite-volume phase shift calculated from Eq. (12). A plot of these estimates, $\delta(k_r)$, for a range of volumes is shown alongside the infinite-volume phase shift as a function of E in Fig. 3. It is clear that the convergence to the infinite-volume phase shift as a function of L is quite slow, with an error of approximately 20 MeV in the resonance position, $E_{\rm res}$, on a 5 fm box (using a linear interpolation between the two energies closest to $E_{\rm res}$).



FIG. 2. (color online). The Breit-Wigner-like phase shift δ associated with elastic $N\pi$-scattering via a Δ-baryon intermediate state, plotted against external energy E, as calculated from the on-shell t-matrix of Eq. (12).



FIG. 3. (color online). Estimates of the phase shift obtained from Lüscher's formula applied to the $\Delta N\pi$ model for a range of L values. The dot-dashed lines that connect the discrete points are merely to guide the eye. The infinite-volume phase shift is also shown as a solid line.



FIG. 4. (color online). The energy differences δE between the exact levels from the $\Delta N\pi$ model and the levels obtained from Lüscher's formula using the exact phase shifts from the t-matrix. Energy level 1 corresponds to the lowest lying state.



FIG. 5. (color online). The resonance energy $E_{\rm res}$ plotted against $1/L$, using a linear and a Breit-Wigner interpolation of the discrete phase shift obtained using Eq. (1). The experimental value is marked with a square. The results from fitting pseudodata with a different regulator, a Gaussian with $\Lambda = 0.5,\ 0.6\ \&\ 0.8$ GeV, are plotted. The results of fitting pseudodata containing a non-resonant background interaction (see Eq. (10)) are also shown, treating the Gaussian scale Λ as a fit parameter.

By applying Lüscher's formula from Eq. (1), the corresponding estimate of the energy levels may be calculated. The energy difference between the exact energy levels from the model and the levels obtained from Lüscher's formula is shown for the five lowest-lying states in Fig. 4. The energy difference is exponentially suppressed for large values of L.

The behaviour of $E_{\rm res}$ as a function of L is of great interest. This provides an indication of the robustness of the method for extracting an estimate of the true phase shift. The result of obtaining $E_{\rm res}$, using a linear and also a Breit-Wigner interpolation, is shown as a function of $1/L$ in Fig. 5. Firstly, there are significant oscillations in the behaviour of $E_{\rm res}$ even up to box sizes of 7 fm. This is a consequence of the behaviour of the geometric ϕ function. Furthermore, the large L behaviour is linear in $1/L$; though it is reasonable to expect that exponential suppression will dominate for larger box sizes than 40 fm. A linear extrapolation of the resonance position in $1/L$ is possible, but Fig. 5 indicates that a reliable result requires a box size of $L \gtrsim 10$ fm. Clearly, these box sizes far exceed feasibility for a current lattice QCD calculation. For presently conceivable lattice sizes ($L \leq 5$ fm), the error in the resonance position obtained through either a linear or a Breit-Wigner interpolation to the point where $\delta = 90°$, yields an error of order 10%. For this reason, we have been led to consider an alternative method for extracting an estimate of the phase shift.

Motivated by the general result that the t-matrix near any resonance is separable [24], the idea of fitting the energy levels from a lattice calculation to a separable model is explored. By treating the coupling constant, χ_Δ, and the bare resonance mass, Δ_0, as fit parameters, the energy levels between the lattice calculation and the previously established $\Delta N\pi$ model can be matched. A new estimate of the exact phase shift can be then calculated from the t-matrix calculation of Eq. (12) using the parameters of the fit.

In order to test this idea, the energy levels for the $\Delta N\pi$ model are generated using a dipole regulator with $\Lambda = 0.8$ GeV, and a value of Δ_0 tuned to reproduce the experimental resonance position. These energy levels are treated as pseudodata. A model is then chosen using a different regulator function, such as a Gaussian regulator. By minimizing the χ^2 function between the two sets of energy levels and obtaining fit values for χ_Δ and Δ_0, exact infinite-volume phase shifts may be calculated for a variety of values of L. Only energy levels in the region where the model is valid, near the resonance ($E_r \lesssim 500$ MeV), are used when matching two sets of energy levels. The resonance position obtained using this procedure is shown in Fig. 5 as a function of $1/L$, for a variety of suitable values of the Gaussian regulator mass. Clearly, refitting the pseudodata with a different regulator function still leads to a result that is stable with respect to L. Furthermore, the extracted resonance position is within a few MeV of the actual resonance position. This is encouraging, because it suggests that approximating the underlying physics of a lattice calculation with a regulator, sensibly chosen, will lead to a result that is not highly dependent on the features of the particular form of the regulator function.

One might argue that the simple model including only $\Delta \leftrightarrow N\pi$ couplings is ideally suited to the reconstruction method proposed here. For that reason, the following case was considered, in which pseudodata were generated from a Hamiltonian model that now includes the effective contact interaction from Eq. (10), again using a dipole regulator with $\Lambda = 0.8$ GeV. The energy levels were fit to the more basic model, with only the $\Delta \leftrightarrow N\pi$ coupling, using a Gaussian regulator, and treating Λ as an additional fit parameter. In this case, the fitted value of the Gaussian regulator is $\Lambda_{\text{fit}} = 0.42$ GeV, leading to a stable extraction of the resonance energy, as shown in Fig. 5. This is remarkable, as it shows that the bare resonance mass, the coefficient of the interaction and the finite-range scale Λ are collectively able to mock-up the missing Chew-Low interaction. That is, more complicated interactions are incorporated into the fit degrees of freedom.

In summary, it has been demonstrated that care must be taken in matching the asymptotic states of elastic scattering to the energy spectrum in a finite volume. The finite range of the hadronic interactions entails a $\sim 1/L$ deviation in the extracted resonance position using Lüscher's method. A new method is proposed in which a $\Delta N\pi$ interaction model is constructed, allowing the bare resonance position and the coupling strength to be free parameters. By matching the energy levels of an *ab initio* finite-volume calculation, such as lattice QCD, with those of the model, the fitted parameters can be used to reconstruct the exact phase shift using the infinite-volume t-matrix calculation. The results show that the resonance position of the new phase shift is both accurate, and relatively independent of L across a wide range of box sizes, including those used in contemporary lattice QCD simulations.

We would like to thank M. Döring and J. Dudek for helpful conversations. This research was supported by the Australian Research Council through the ARC Centre of Excellence for Particle Physics at the Terascale, and through grants DP110101265 (DBL & RDY) and FL0992247 (AWT).

[1] R. G. Edwards, J. J. Dudek, D. G. Richards, and S. J. Wallace, Phys.Rev. **D84**, 074508 (2011), 1104.5152.

[2] G. P. Engel, C. Lang, M. Limmer, D. Mohler, and A. Schafer (BGR [Bern-Graz-Regensburg] Collaboration), Phys.Rev. **D82**, 034505 (2010), 1005.1748.

[3] M. S. Mahbub, W. Kamleh, D. B. Leinweber, P. J. Moran, and A. G. Williams (CSSM Lattice collaboration), Phys.Lett. **B707**, 389 (2012), 1011.5724.

[4] S. He, X. Feng, and C. Liu, JHEP **0507**, 011 (2005), hep-lat/0504019.

[5] V. Bernard, M. Lage, U.-G. Meißner, and A. Rusetsky, JHEP **0808**, 024 (2008), 0806.4495.

[6] V. Bernard, D. Hoja, U.-G. Meißner, and A. Rusetsky, JHEP **0906**, 061 (2009), 0902.2346.

[7] E. Oset, M. Döring, U.-G. Meißner, and A. Rusetsky (2011), 1108.3923.

[8] M. Döring and U.-G. Meißner, JHEP **1201**, 009 (2012), 1111.0616.

[9] M. Döring, U.-G. Meißner, E. Oset, and A. Rusetsky, Eur.Phys.J. **A47**, 139 (2011), 1107.3988.

[10] L. Roca and E. Oset, Phys.Rev. **D85**, 054507 (2012), 1201.0438.

[11] P. Giudice, D. McManus, and M. Peardon (2012), 1204.2745.

[12] S. Kreuzer and H. W. Grießhammer (2012), 1205.0277.

[13] M. Albaladejo, J. Oller, E. Oset, G. Rios, and L. Roca (2012), 1205.3582.

[14] M. Gockeler, R. Horsley, M. Lage, U.-G. Meißner, P. Rakow, et al. (2012), 1206.4141.

[15] M. Luscher, Nucl.Phys. **B354**, 531 (1991).

[16] R. D. Young, D. B. Leinweber, and A. W. Thomas, Prog. Part. Nucl. Phys. **50**, 399 (2003), hep-lat/0212031.

[17] D. B. Leinweber, A. W. Thomas, and R. D. Young, Phys. Rev. Lett. **92**, 242002 (2004), hep-lat/0302020.

[18] D. B. Leinweber, A. W. Thomas, and R. D. Young, Nucl.Phys. **A755**, 59 (2005), hep-lat/0501028.

[19] J. Hall, D. Leinweber, and R. Young, Phys.Rev. **D82**, 034010 (2010), 1002.4924.

[20] J. Hall, F. Lee, D. Leinweber, K. Liu, N. Mathur, et al., Phys.Rev. **D84**, 114011 (2011), 1101.4411.

[21] S. Theberge, A. W. Thomas, and G. A. Miller, Phys.Rev. **D22**, 2838 (1980).

[22] A. W. Thomas, S. Theberge, and G. A. Miller, Phys.Rev. **D24**, 216 (1981).

[23] T. Mongan, Phys.Rev. **178**, 1597 (1969).

[24] C. Lovelace, Phys.Rev. **135**, B1225 (1964).